

## MDR-LIKE ABC TRANSPORTER GENE FROM PLANTS

This application is a continuation in part of International Application No. PCT/US99/22363, filed September 24, 1999, which claims priority under 35 U.S.C. §120 to U.S. Provisional Application 60/101,814, the entireties of both of which are incorporated by reference herein.

Pursuant to 35 U.S.C. §202(c), it is acknowledged that the U.S. Government has certain rights in the invention described herein, which was made in part with funds from the National Science Foundation, Grant No. IBN-9416016.

### FIELD OF THE INVENTION

This invention relates to the field of stress resistance in plants. In particular, the invention provides a novel gene from plants, which encodes an MDR-like ABC transporter, involved in detoxification of certain xenobiotics to protect plants from their detrimental effects.

### BACKGROUND OF THE INVENTION

Several publications are referenced in this application to describe the state of the art to which the invention pertains. Each of these publications is incorporated by reference herein.

Environmental stress is one of the most important limitations on plant productivity, growth and survival. An ever-increasing source of environmental stress to plants is

the stress caused by environmental pollutants in the soil, water and atmosphere. Such pollutants include herbicides, pesticides and related agronomic products, as well as organic and inorganic waste material from industry and other sources.

- 5 Other toxic agents that threaten the survival of plants include various toxins produced by ephiphytic or soilborne microorganisms, such as fungi and bacteria.

In order to survive in toxic environments, plants must have mechanisms to detoxify xenobiotics, heavy metals  
10 and other toxic compounds. This generally involves modification of the toxic compound and subsequent excretion into the vacuole or apoplastic space. Recently, certain ATP-binding cassette (ABC) transporters have been identified in plants, which appear to be involved in the detoxification  
15 process.

The ABC transporter family is very large, with representatives existing in many different classes of organisms. Two well studied groups of ABC transporters, encoded by *mdr* and *mrp* genes, respectively, are associated  
20 with the multi-drug resistance phenomenon observed in mammalian tumor cells. The *mdr* genes encode a family of P-glycoproteins that mediate the energy-dependent efflux of certain lipophilic drugs from cells. The *mrp* genes encode a family of transporters that mediate the extrusion of a  
25 variety of organic compounds after their conjugation with glutathione. *YCF1*, the yeast homolog of *mrp*, encodes a protein capable of glutathione-mediated detoxification of heavy metals.

Homologs of *mrp* and *mdr* genes have been identified  
30 in plant species. In *Arabidopsis thaliana*, the glutathione-

conjugate transporter encoded by the *mrp* homolog is located in the vacuolar membrane and is responsible for sequestration of xenobiotics in the central vacuole (Tommasini et al., FEBS Lett. 411: 206-210, 1997; Li et al., Plant Physiol. 107: 1257-1268, 1995). An *mdr*-like gene (*atpgp1*) has also been identified in *A. thaliana*, which encodes a putative P-glycoprotein homolog. The *atpgp1* gene was found to share significant sequence homology and structural organization with human *mdr* genes, and was expressed with particular abundance in inflorescence axes (Dudler & Hertig, J. Biol. Chem. 267: 5882-5888, 1992). Other MDR homologs have been found in potato (Wang et al., Plant Mol. Biol. 31: 683, 1996) and barley (Davies et al., Gene 199: 195, 1997).

The aforementioned *mrp* and *mdr* plant homologs were identified as a result of an effort to understand the molecular basis for development in plants of cross-resistance to herbicides of unrelated classes. However, these transporters are likely to serve the more general role in plants of sequestering, secreting, or otherwise detoxifying various organic and inorganic xenobiotics. Accordingly, it will constitute an advance in the art of plant genetic engineering of stress tolerance to identify and characterize other members of this class of transporters in plants.

## SUMMARY OF THE INVENTION

In accordance with the present invention, a new plant *mdr* homolog, referred to herein as *plPAC*, has been identified. Unlike the previously identified plant *mdr* homologs, this new gene is inducible by a class of compounds known to inhibit chloride ion channels. The new gene is also

inducible by auxin and binds NPA.

According to one aspect of the invention, a nucleic acid isolated from a plant is provided, which encodes a p-glycoprotein that is inducible by exposure of the plant to NPPB or auxin. The isolated nucleic acid is preferentially expressed in plant roots upon exposure of the plant to NPPB. In a preferred embodiment, the plant from which the nucleic acid is isolated is selected from the group consisting of *Brassica napus* and *Arabidopsis thaliana* and is 3850-4150 nucleotides in length. In a more preferred embodiment, the nucleic acid has the restriction sites shown in Figure 4 for at least three restriction enzymes. In particularly preferred embodiments, the nucleic acid molecule encodes a polypeptide having SEQ ID NO:2. In an exemplary embodiment, the nucleic acid is a cDNA comprising the coding region of SEQ ID NO:1 or SEQ ID NO:10.

According to another aspect of the invention is an expression cassette that comprises a plPAC gene operably linked to a promoter, and in a more preferred embodiment the plPAC gene is from *Arabidopsis*. In preferred embodiments, the expression cassette comprises the cauliflower mosaic virus 35S promoter, and part of all of SEQ ID NO:1 or SEQ ID NO:10. Further included in this aspect is a vector comprising the expression cassette and a method for producing transgenic plants with the expression cassette and vector.

Another aspect of the invention is drawn to transgenic cells and plants containing the nucleic acids of the invention. In one preferred embodiment, the nucleic acids are be in the aforementioned expression cassette.

Further included in this aspect are reproductive units from

the transgenic plant.

According to another aspect of the invention, an isolated nucleic acid molecule is provided, which has a sequence selected from the group consisting of: a) SEQ ID NO:1 and SEQ ID NO:10; b) a nucleic acid sequence that is at least about 60% homologous to the coding regions of SEQ ID NO:1 or SEQ ID NO:10; c) a sequence hybridizing with SEQ ID NO:1 or SEQ ID NO:10 at moderate stringency; d) a sequence encoding part or all of a polypeptide having SEQ ID NO:2; e) a sequence encoding an amino acid sequence that is at least about 70% identical to SEQ ID NO:2; f) a sequence encoding an amino acid sequence that is at least about 80% similar to SEQ ID NO:2; g) a sequence encoding an amino acid sequence that is at least about 40% similar to residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2; and h) a sequence hybridizing at moderate stringency to a sequence encoding residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2. A polypeptide produced by expression of the above listed sequences is also provided.

According to another aspect of the invention, an isolated plant p-glycoprotein, which is inducible upon exposure of the plant to NPPB, is provided. The polypeptide preferably confers upon a cell in which it is found resistance to Rhodamine 6G. The polypeptide is preferentially produced in roots upon the exposure to the NPPB. The polypeptide is preferably from *Brassica napus* or *Arabidopsis thaliana*. In most preferred embodiments, the polypeptide has a sequence that is a) an amino acid sequence that is at least 80% similar to SEQ ID NO:2; b) an amino acid sequence that is at least 70% identical to SEQ ID NO:2; c) an amino acid sequence that is at least 40% similar to residues

1-76, 613-669 or 1144-1161 of SEQ ID NO:2; and d) an amino acid sequence encoded by a nucleic acid sequence hybridizing at moderate stringency to a amino acid sequence encoding residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2.

5           According to other aspects of the invention, antibodies immunologically specific for the polypeptides of the invention are provided, that immunologically specific to any of the polypeptides, of polypeptide encoded by the nucleic acids of the invention. In a preferred embodiment, 10 the antibody is immunospecific to residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2.

          According to another aspect of the invention, a plant p-glycoprotein gene promoter, which is inducible by NPPB, is also provided. In a preferred embodiment, the 15 promoter is part or all of residues 1-3429 of SEQ ID NO:10. According to another aspect of the invention, plants that have reduces levels of *plPAC* protein are provided. In a preferred embodiment, these plants have mutations in the *plPAC* gene, and in a particularly preferred embodiment, the 20 *plPAC* gene is mutated due to the insertion of a T-DNA. Also provided with this aspect is a method for selecting plants with mutations in *plPAC* using SEQ ID NOS:11-14 as PCR primers.

          These and other features and advantages of the 25 present invention will be described in greater detail in the description and examples set forth below.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

**Figure 1.** Amino acid sequence lineup of *ATPAC* 30 deduced amino acid sequence and the amino acid sequences of

related mammalian and plant genes. The lineup shows the ATPAC deduced amino acid sequence (SEQ ID NO:2) compared with (1) hmdr1 (SEQ ID NO:3); (2) mmdr1 (SEQ ID NO: 4); (3) hmdr3 (SEQ ID NO:5); (4) mmdr2 (SEQ ID NO:6); (5) atpgp1 (SEQ ID NO:7); and (6) atpgp2 (SEQ ID NO:8). A consensus sequence (SEQ ID NO: 9) is also shown.

**Figure 2.** Graph depicting the effect of rhodamine 6G on the growth rate of cells transformed with and expressing ATPAC as compared with control cells not containing ATPAC.

**Figure 3.** Restriction map of genomic clone of ATPAC, SEQ ID NO:10.

**Figure 4.** Restriction map of cDNA clone of ATPAC, SEQ ID NO:1.

## DETAILED DESCRIPTION OF THE INVENTION

### I. Definitions

Various terms relating to the biological molecules of the present invention are used hereinabove and also throughout the specification and claims.

With reference to nucleic acids of the invention, the term "isolated nucleic acid" is sometimes used. This term, when applied to DNA, refers to a DNA molecule that is separated from sequences with which it is immediately contiguous (in the 5' and 3' directions) in the naturally occurring genome of the organism from which it was derived. For example, the "isolated nucleic acid" may comprise a DNA molecule inserted into a vector, such as a plasmid or virus vector, or integrated into the genomic DNA of a procaryote or eucaryote. An "isolated nucleic acid molecule" may also

comprise a cDNA molecule.

With respect to RNA molecules of the invention the term "isolated nucleic acid" primarily refers to an RNA molecule encoded by an isolated DNA molecule as defined  
5 above. Alternatively, the term may refer to an RNA molecule that has been sufficiently separated from RNA molecules with which it would be associated in its natural state (i.e., in cells or tissues), such that it exists in a "substantially pure" form (the term "substantially pure" is defined below).

10 Nucleic acid sequences and amino acid sequences can be compared using computer programs that align the similar sequences of the nucleic or amino acids thus define the differences. For purposes of this invention, the DNASTar program (DNASTar, Inc., Madison, Wisconsin) and the default  
15 parameters used by that program are the parameters intended to be used herein to compare sequence identity and similarity. Alternately, the Blastn and Blastp 2.0 programs provided by the National Center for Biotechnology Information (at <http://www.ncbi.nlm.nih.gov/blast/> ; Altschul et al.,  
20 1990, J Mol Biol 215:403410) using a gapped alignment with default parameters, may be used to determine the level of identity and similarity between nucleic acid sequences and amino acid sequences.

The term "substantially the same" refers to nucleic  
25 acid or amino acid sequences having sequence variation that do not materially affect the nature of the protein (i.e. the structure, thermostability characteristics and/or biological activity of the protein). With particular reference to nucleic acid sequences, the term "substantially the same" is  
30 intended to refer to the coding region and to conserved



sequences governing expression, and refers primarily to degenerate codons encoding the same amino acid, or alternate codons encoding conservative substitute amino acids in the encoded polypeptide. With reference to amino acid sequences, the term "substantially the same" refers generally to conservative substitutions and/or variations in regions of the polypeptide not involved in determination of structure or function.

The terms "percent identical" and "percent similar" are also used herein in comparisons among amino acid and nucleic acid sequences. When referring to amino acid sequences, "percent identical" refers to the percent of the amino acids of the subject amino acid sequence that have been matched to identical amino acids in the compared amino acid sequence by a sequence analysis program. "Percent similar" refers to the percent of the amino acids of the subject amino acid sequence that have been matched to identical or conserved amino acids. Conserved amino acids are those which differ in structure but are similar in physical properties such that the exchange of one for another would not appreciably change the tertiary structure of the resulting protein. Conservative substitutions are defined in Taylor (1986, J. Theor. Biol. 119:205). When referring to nucleic acid molecules, "percent identical" refers to the percent of the nucleotides of the subject nucleic acid sequence that have been matched to identical nucleotides by a sequence analysis program.

With respect to protein, the term "isolated protein" or "isolated and purified protein" is sometimes used herein. This term refers primarily to a protein produced by

expression of an isolated nucleic acid molecule of the invention. Alternatively, this term may refer to a protein which has been sufficiently separated from other proteins with which it would naturally be associated, so as to exist  
5 in "substantially pure" form.

The term "substantially pure" refers to a preparation comprising at least 50-60% by weight the compound of interest (e.g., nucleic acid, oligonucleotide, protein, etc.). More preferably, the preparation comprises at least  
10 75% by weight, and most preferably 90-99% by weight, the compound of interest. Purity is measured by methods appropriate for the compound of interest (e.g. chromatographic methods, agarose or polyacrylamide gel electrophoresis, HPLC analysis, and the like).

15 With respect to antibodies of the invention, the term "immunologically specific" refers to antibodies that bind to one or more epitopes of a protein of interest, but which do not substantially recognize and bind other molecules in a sample containing a mixed population of antigenic  
20 biological molecules.

With respect to oligonucleotides, the term "specifically hybridizing" refers to the association between two single-stranded nucleotide molecules of sufficiently complementary sequence to permit such hybridization under  
25 pre-determined conditions generally used in the art (sometimes termed "substantially complementary"). In particular, the term refers to hybridization of an oligonucleotide with a substantially complementary sequence contained within a single-stranded DNA or RNA molecule of the  
30 invention, to the substantial exclusion of hybridization of

the oligonucleotide with single-stranded nucleic acids of non-complementary sequence.

5 The term "expression cassette", as used herein, comprises 5' and 3' regulatory regions operably linked to a coding sequence. The coding sequence may be in the sense or antisense orientation with respect to the 5' regulatory region.

The term "promoter region" refers to the 5' regulatory regions of a gene.

10 The term "reporter gene" refers to genetic sequences which may be operably linked to a promoter region forming a transgene, such that expression of the reporter gene coding region is regulated by the promoter and expression of the transgene is readily assayed.

15 The term "selectable marker gene" refers to a gene product that when expressed confers a selectable phenotype, such as antibiotic resistance, on a transformed cell or plant.

20 The term "operably linked" means that the regulatory sequences necessary for expression of the coding sequence are placed in the DNA molecule in the appropriate positions relative to the coding sequence so as to effect expression of the coding sequence. This same definition is sometimes applied to the arrangement of coding sequences and transcription control elements (e.g. promoters, enhancers, and termination elements) in an expression vector.

25 The term "DNA construct" refers to genetic sequence used to transform plants and generate progeny transgenic plants. These constructs may be administered to plants in a viral or plasmid vector. Other methods of delivery such as

30

Agrobacterium T-DNA mediated transformation and transformation using the biolistic process are also contemplated to be within the scope of the present invention. The transforming DNA may be prepared according to standard protocols such as those set forth in "Current Protocols in Molecular Biology", eds. Frederick M. Ausubel et al., John Wiley & Sons, 2001.

The term "xenobiotic" refers to foreign chemicals or agents not produced or naturally found in the organism. The term is commonly used in reference to toxic or otherwise detrimental foreign chemicals, such as organic pollutants or heavy metals.

## II. Description of *plPAC* and its Encoded Polypeptide

In accordance with the present invention, a nucleic acid encoding a novel ATP-binding-cassette (ABC) transporter has been isolated and cloned from plants. This novel ABC transporter is induced by auxin and binds NPA. The nucleic acid is referred to herein as *plPAC*.

A cDNA clone of the *plPAC* from *Arabidopsis thaliana*, an exemplary *plPAC* of the invention, is described in detail herein and its nucleotide sequence is set forth in Example 1 as SEQ ID NO:1. This nucleic acid molecule is referred to as "ATPAC". It is 36% identical and 51% similar to human *mdr1* across the entire sequence. It is 51% identical to the *atpgp1* gene reported by Dudler & Hertig (1997, *supra*) and 50% identical to *atpgp2*, a close homolog of *atpgp1*, published in the Genbank database. ATPAC protein is 65% similar to *atpgp1* and *atpgp2* proteins.

A partial clone of a *plPAC* of the invention was originally isolated from *Brassica napus* via differential expression screening of plants grown in the presence or absence of the chloride channel blocker, 5-nitro-2-(3-phenylpropylamino) benzoic acid (NPPB). A 0.5 kb gene fragment was identified, which had been up-regulated in response to NPPB treatment. This cDNA fragment was used to screen an *Arabidopsis* cDNA library, from which the complete *ATPAC* clone was isolated. The isolation and characterization of *ATPAC* is described in Example 1.

A genomic clone of *ATPAC* (SEQ ID NO:10) has also been isolated from a bacterial artificial chromosome (BAC) library of the *Arabidopsis* genome (BAC clone IGF F3J22, obtained from the *Arabidopsis* stock center, Ohio State University). A 7 kb fragment containing part of *ATPAC* and additional 5' regulatory sequences was subcloned into a plasmid vector (pBluescript). A restriction map of *ATPAC* is found in Fig. 3. The corresponding cDNA clone of *ATPAC* is found in SEQ ID NO:1 and its restriction map is Fig. 4.

In *Arabidopsis*, *plPAC* of the present invention is expressed in roots, leaves, flowers, and shoot meristem. Expression is strong in the hypocotyl of etiolated seedlings, but not in light grown seedlings. Expression of *ATPAC* is also relatively high in cotyledons, meristem, roots and the first true leaves of seedlings.

Among the unique features of this nucleic acid molecule as compared with other *mdr*-like genes from plants are its inducibility by certain compounds, including NPPB, herbicides and auxin, and its preferential expression in roots. The promoter regulatory region of *ATPAC* comprises

residues 1-3429 of SEQ ID NO:10.

Although the *ATPAC* cDNA clone from *Arabidopsis thaliana* is described and exemplified herein, this invention is intended to encompass nucleic acid sequences and proteins from other plant species that are sufficiently similar to be used instead of *ATPAC* nucleic acid and proteins for the purposes described below. These include, but are not limited to, allelic variants and natural mutants of SEQ ID NO:1, which are likely to be found in different species of plants or varieties of *Arabidopsis*.

Because such variants are expected to possess certain differences in nucleotide and amino acid sequence, this invention provides an isolated *plPAC* nucleic acid molecule having at least about 60% (preferably 70% and more preferably over 80%) sequence homology in the coding regions with the nucleotide sequence set forth as SEQ ID NO:1 or SEQ ID NO:10 (and, most preferably, specifically comprising the coding region of SEQ ID NO:1). Also provided are nucleic acids that encode a polypeptide that is at least about 40% (preferably 50% and most preferably 60%) similar to residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2. Also provided are nucleic acids that hybridize to the nucleic acids of SEQ ID NO:1, SEQ ID NO:10, or nucleic acids encoding the regions of residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2, preferably under moderate stringency (more preferably, high stringency, and most preferably, very high stringency).

In other preferred embodiments, the nucleic acids have a restriction digest map that is identical for at least 3 enzymes (more preferably 6 enzymes and most preferably 9 enzymes) to the maps shown in Figs. 3 or 4. In another

preferred embodiment, the nucleic acids have a restriction digest map identical to those shown in Fig. 3 for enzymes *XhoI*, *XcmI* and *SpeI* (preferably additionally *SacI*, *PacI* and *BsaI*, and most preferably additionally *AclI*, *BanI* and *SnaBI*).

5 In another preferred embodiment, the nucleic acids have a restriction digest map identical to those shown in Fig. 4 for enzymes *XbaI*, *TatI* and *NciI* (preferably additionally *DraI*, *BsmI* and *BclI*, and most preferably additionally *AccI*, *BsgI* and *TliI*). The nucleic acids of the invention are at least 20 nucleic acids in length (preferably at least 50 nucleic acids and most preferably at least 100 nucleic acids).

10 In accordance with the invention, novel *plPAC* genes from two plant species, *Brassica napus* and *Arabidopsis thaliana*, are presented. This constitutes the first description of this unique p-glycoprotein in plants. Indeed, the closest known protein sequence, also from *Arabidopsis*, is only 65% identical suggesting that the *ATPAC* gene is novel and is expected to have novel properties. The isolation of two *plPAC* genes from different species enables the isolation of further *plPAC* genes from other plant species. Isolated nucleic acids that are *plPAC* genes from any plant species are considered part of the instant invention. In particular, the nucleic acids of other *plPAC* genes can be isolated using sequences of *ATPAC* that distinguish *plPAC* genes from other plant *mdr* genes according to methods that are well known to those in the art of gene isolation. In particular, sequences that encode residues 1-76, 613-669 and 1144-1161 of SEQ ID NO:2 can be used. In a preferred embodiment, the *plPAC* gene is from any higher plant species (more preferred from a dicot

15  
20  
25  
30

species, and most preferred from a species in Brassicaceae (or Cruciferae)).

This invention also provides isolated polypeptide products of the open reading frames of SEQ ID NO:1 or SEQ ID NO:10, having at least about 70% (preferably 80% and most preferably 90%) sequence identity, or at least about 80% similarity (preferably 90% and more preferably 95%) with the amino acid sequence of SEQ ID NO:2. In another embodiment, the polypeptides of the invention are at least about 40% identical (preferably 50%, and most preferably 60%) to the regions of residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2. Because of the natural sequence variation likely to exist among *plPAC* genes, one skilled in the art would expect to find up to about 30-40% nucleotide sequence variation, while still maintaining the unique properties of the *plPAC* gene and encoded polypeptide of the present invention. Such an expectation is due in part to the degeneracy of the genetic code, as well as to the known evolutionary success of conservative amino acid sequence variations, which do not appreciably alter the nature of the encoded protein. Accordingly, such variants are considered substantially the same as one another and are included within the scope of the present invention.

Also provided are transgenic plants transformed with part or all of the nucleic acids of the invention. Transgenic plants that over-express a *plPAC* coding sequence are one embodiment of this aspect of the invention. Example 3 provides for one prototype of such a plant. In a preferred embodiment, the *ATPAC* gene is used, and in a most preferred embodiment SEQ ID NO:1 or SEQ ID NO:10 is used. The *plPAC*



gene may be placed under a powerful constitutive promoter, such as the Cauliflower Mosaic Virus (CaMV) 35S promoter or the figwort mosaic virus 35S promoter. In a preferred embodiment, the 35SCaMV promoter is used. Transgenic plants  
5 expressing the *plPAC* gene under an inducible promoter (either its own promoter or a heterologous promoter) are also contemplated to be within the scope of the present invention. Inducible plant promoters include the tetracycline repressor/operator controlled promoter. In a preferred  
10 embodiment, a native *plPAC* promoter is used, and in a most preferred embodiment, residues 1-3429 of SEQ ID NO:10 is used. Plant species that are contemplated for overexpression of a *plPAC* coding sequence include, but are not limited to, soybean.

15 In another embodiment, overexpression of *plPAC* is induced to generate a co-suppression effect. This excess expression serves to promote down-regulation of both endogenous and exogenous *plPAC* genes.

In some instances, it may be desirable to down-  
20 regulate or inhibit expression of endogenous *plPAC* in plants possessing the gene. Accordingly, *plPAC* nucleic acid molecules, or fragments thereof, may also be utilized to control the production of *plPAC*-encoded P-glycoproteins. In one embodiment, full-length *plPAC* antisense molecules or  
25 antisense oligonucleotides, targeted to specific regions of *plPAC*-encoded RNA that are critical for translation, are used. The use of antisense molecules to decrease expression levels of a pre-determined gene is known in the art. In a preferred embodiment, antisense molecules are provided in  
30 *situ* by transforming plant cells with a DNA construct which,

upon transcription, produces the antisense sequences. Such constructs can be designed to produce full-length or partial antisense sequences. One example of antisense *plPAC* transgenic plants is given in Example 3.

5 In another embodiment, knock-out plants are obtained by screening a T-DNA mutagenized plant population for insertions in the *plPAC* gene (see Krysan et al., 1996, PNAS 93:8145). One example of this embodiment of the invention is found in Example 3. Optionally, transgenic  
10 plants can be created containing mutations in the region encoding the active site of *plPAC*. These last two embodiments are preferred over the use of anti-sense constructs due to the high homology among P-glycoproteins. The promoter of *ATPAC* is also provided in accordance with the  
15 invention. This promoter has the useful properties of root expression and inducability by NPPB. Presence of NPPB in the growth medium of *Arabidopsis* seedlings results in increased expression of *ATPAC* of the present invention.

Further, when approximately 4kb of upstream *ATPAC*  
20 promoter DNA is fused to the GUS reporter gene and transformed into wild-type plants, GUS staining is strong in the hypocotyl of etiolated seedlings, but not in light grown seedlings. Further, expression is high in cotyledons, meristem, root, and the first true leaves of seedlings.  
25 Staining was also observed in flowers and the apical portion of the inflorescence.

The prototypic example of this aspect of the invention is residues 1-3429 of SEQ ID NO:10. It is anticipated that *plPAC* genes from other plant species will  
30 likewise exhibit the aforementioned useful properties. As

these promoter regions can easily be isolated from the *plPAC* genes that are provided with the invention, all plant *plPAC* gene promoters are provided with the invention. The nucleic acids of the invention therefore include a nucleic acid molecule that is at least about 70% identical (preferably 80% and most preferably 90%) to the residues 1-3429 of SEQ ID NO:10. Also provided are nucleic acids that hybridize to the nucleic acid residues 1-3429 of SEQ ID NO:10 preferably under moderate stringency (more preferably, high stringency, and most preferably, very high stringency).

Thus, the *PlPAC* of the present invention encodes an ABC transporter that binds NPA and is involved with auxin transport in the plant. Mutants of *Arabidopsis* lacking *ATPAC* and double mutants lacking both *ATPAC* and *AtPGP1* display morphological phenotypes consistent with their demonstrated impairments in polar auxin transport. It has been widely accepted that NPA-sensitive regulatory site and the auxin-conducting channel of the efflux carrier are separate molecular entities. Strong evidence indicates that PIN-like genes encode the auxin-conducting channel of the efflux carrier (Palme and Galweiler (1999) Curr. Op. Plant Biol. 2:375-381). However, as evidenced in Example 4 of the present invention, MDR-like genes are components of the NPA-sensitive regulatory site.

Expression of the *plPAC* gene of the present invention is inducible by auxin as described below in Example 4. Treatment of wild type *Arabidopsis* seedlings with 2,4-D or high concentrations of auxin result in plants with the same phenotype as that of *ATPAC* mutant plants. The strongly curved organs of the double mutant resemble the twisted forms

induced by treatment of some dicots with herbicidal levels of the auxin analog, 2,4-D and in *Arabidopsis* by treatment with auxin transport inhibitors (Sieburth, (1999) Plant Physiol. 121: 1179-1190). This indicates that ATPAC of the present invention pumps auxin or auxin conjugates from sites of synthesis, such as the apical meristem and expanding cotyledons (Sachs, (1991) Development 117: 833-893). Under this model, tissues that express this pump would accumulate auxin as a result of the mutation and the altered auxin balance could be responsible for the altered growth patterns typifying the ATPAC phenotype. Further support of this model is the similarity of auxin to indolylic substrates pumped by human MDR1, and the finding that ATPAC expression is increased by auxin. Also, the fact that NPA binds to ATPAC and that *atpac* knockout mutants can be phenocopied by auxin application suggests that ATPAC is an important component of the auxin transport and distribution machinery.

The present invention also provides antibodies capable of immunospecifically binding to polypeptides of the invention. In a preferred embodiment, the antibodies react immunospecifically with various epitopes of the *plPAC*-encoded polypeptides. In a particularly preferred embodiment, the antibodies are immunologically specific to the polypeptide of residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2.

The following sections sets forth the general procedures involved in practicing the present invention. To the extent that specific materials are mentioned, it is merely for purposes of illustration and is not intended to limit the invention. Unless otherwise specified, general cloning procedures, such as those set forth in Sambrook et

al., Molecular Cloning, Cold Spring Harbor Laboratory (1989)  
(hereinafter "Sambrook et al.") or Ausubel et al. (eds)  
Current Protocols in Molecular Biology, John Wiley & Sons  
(2001) (hereinafter "Ausubel et al.") are used.

5

**III. Preparation of *PlPAC* Nucleic Acid Molecules,  
encoded Polypeptides, Antibodies Specific for the  
Polypeptides and Transgenic Plants**

10

**1. Nucleic Acid Molecules**

15

*PlPAC* nucleic acid molecules of the invention may  
be prepared by two general methods: (1) they may be  
synthesized from appropriate nucleotide triphosphates, or (2)  
they may be isolated from biological sources. Both methods  
utilize protocols well known in the art.

20

25

30

The availability of nucleotide sequence  
information, such as the cDNA having SEQ ID NO:1, enables  
preparation of an isolated nucleic acid molecule of the  
invention by oligonucleotide synthesis. Synthetic  
oligonucleotides may be prepared by the phosphoramidite  
method employed in the Applied Biosystems 38A DNA Synthesizer  
or similar devices. The resultant construct may be purified  
according to methods known in the art, such as high  
performance liquid chromatography (HPLC). Long, double-  
stranded polynucleotides, such as a DNA molecule of the  
present invention, must be synthesized in stages, due to the  
size limitations inherent in current oligonucleotide  
synthetic methods. Thus, for example, a long double-stranded  
molecule may be synthesized as several smaller segments of  
appropriate complementarity. Complementary segments thus  
produced may be annealed such that each segment possesses

appropriate cohesive termini for attachment of an adjacent segment. Adjacent segments may be ligated by annealing cohesive termini in the presence of DNA ligase to construct an entire long double-stranded molecule. A synthetic DNA  
5 molecule so constructed may then be cloned and amplified in an appropriate vector.

*PlPAC* genes also may be isolated from appropriate biological sources using methods known in the art. In fact, the *ATPAC* clone was isolated from an *Arabidopsis* cDNA library  
10 using a partial clone obtained from *Brassica napus*. In alternative embodiments, genomic clones of *plPAC* may be isolated.

In accordance with the present invention, nucleic acids having the appropriate level sequence homology with  
15 part or all the coding regions of SEQ ID NO:1 or SEQ ID NO:10 may be identified by using hybridization and washing conditions of appropriate stringency. For example, hybridizations may be performed, according to the method of Sambrook et al., using a hybridization solution comprising:  
20 5X SSC, 5X Denhardt's reagent, 1.0% SDS, 100  $\mu$ g/ml denatured, fragmented salmon sperm DNA, 0.05% sodium pyrophosphate and up to 50% formamide. Hybridization is carried out at 37-42°C for at least six hours. Following hybridization, filters are washed as follows: (1) 5 minutes  
25 at room temperature in 2X SSC and 1% SDS; (2) 15 minutes at room temperature in 2X SSC and 0.1% SDS; (3) 30 minutes-1 hour at 37°C in 2X SSC and 0.1% SDS; (4) 2 hours at 45-55° in 2X SSC and 0.1% SDS, changing the solution every 30 minutes. One common formula for calculating the stringency conditions  
30 required to achieve hybridization between nucleic acid

molecules of a specified sequence homology (Sambrook et al., 1989):

$$T_m = 81.5^{\circ}\text{C} + 16.6\text{Log} [\text{Na}^+] + 0.41(\% \text{ G+C}) - 0.63 (\% \text{ formamide}) - 600/\text{\#bp in duplex}$$

5

As an illustration of the above formula, using  $[\text{Na}^+] = [0.368]$  and 50% formamide, with GC content of 42% and an average probe size of 200 bases, the  $T_m$  is  $57^{\circ}\text{C}$ . The  $T_m$  of a DNA duplex decreases by 1 -  $1.5^{\circ}\text{C}$  with every 1% decrease in homology. Thus, targets with greater than about 75% sequence identity would be observed using a hybridization temperature of  $42^{\circ}\text{C}$ .

10

The stringency of the hybridization and wash depend primarily on the salt concentration and temperature of the solutions. In general, to maximize the rate of annealing of the probe with its target, the hybridization is usually carried out at salt and temperature conditions that are 20- $25^{\circ}\text{C}$  below the calculated  $T_m$  of the of the hybrid. Wash conditions should be as stringent as possible for the degree of identity of the probe for the target. In general, wash conditions are selected to be approximately 12- $20^{\circ}\text{C}$  below the  $T_m$  of the hybrid. In regards to the nucleic acids of the current invention, a moderate stringency hybridization is defined as hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100  $\mu\text{g/ml}$  denatured salmon sperm DNA at  $42^{\circ}\text{C}$ , and wash in 2X SSC and 0.5% SDS at  $55^{\circ}\text{C}$  for 15 minutes. A high stringency hybridization is defined as hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100  $\mu\text{g/ml}$  denatured salmon sperm DNA at  $42^{\circ}\text{C}$ , and wash in 1X SSC and

15

20

25

0.5% SDS at 65°C for 15 minutes. A very high stringency hybridization is defined as hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100  $\mu$ g/ml denatured salmon sperm DNA at 42°C, and wash in 0.1X SSC and 0.5% SDS at 65°C for 15 minutes.

Nucleic acids of the present invention may be maintained as DNA in any convenient cloning vector. In a preferred embodiment, clones are maintained in plasmid cloning/expression vector, such as pGEM-T (Promega Biotech, Madison, WI) or pBluescript (Stratagene, La Jolla, CA), either of which is propagated in a suitable *E. coli* host cell.

*plPAC* nucleic acid molecules of the invention include cDNA, genomic DNA, RNA, and fragments thereof which may be single- or double-stranded. Thus, this invention provides oligonucleotides (sense or antisense strands of DNA or RNA) having sequences capable of hybridizing with at least one sequence of a nucleic acid molecule of the present invention, such as selected segments of SEQ ID NO:1 or SEQ ID NO:10. Such oligonucleotides are useful as probes for detecting *plPAC* genes or mRNA in test samples, e.g. by PCR amplification, mapping of genes or for the positive or negative regulation of expression of *plPAC* genes at or before translation of the mRNA into proteins.

The *plPAC* promoter is also expected to be useful in connection with the present invention, inasmuch as it is inducible in plants upon exposure to anion channel blockers. As mentioned above, seven-kilobase fragment of genomic DNA has been isolated, which contains part or all of the *plPAC* promoter from *Arabidopsis thaliana*. This promoter can be



used in chimeric gene constructs to facilitate inducible expression of any coding sequence of interest, upon exposure to NPPB or similar-acting compounds.

5                    **2. Proteins and Antibodies**

Polypeptides encoded by *plPAC* nucleic acids of the invention may be prepared in a variety of ways, according to known methods. If produced *in situ* the polypeptides may be purified from appropriate sources, e.g., plant roots or other  
10 plant parts.

Alternatively, the availability of nucleic acid molecules encoding the polypeptides enables production of the proteins using *in vitro* expression methods known in the art. For example, a cDNA or gene may be cloned into an appropriate  
15 *in vitro* transcription vector, such a pSP64 or pSP65 for *in vitro* transcription, followed by cell-free translation in a suitable cell-free translation system, such as wheat germ or rabbit reticulocytes. *In vitro* transcription and translation systems are commercially available, e.g., from Promega  
20 Biotech, Madison, Wisconsin or BRL, Rockville, Maryland. According to a preferred embodiment, larger quantities of *plPAC*-encoded polypeptide may be produced by expression in a suitable procaryotic or eucaryotic system. For example, part or all of a DNA molecule, such as the cDNA having SEQ ID  
25 NO:1, may be inserted into a plasmid vector adapted for expression in a bacterial cell (such as *E. coli*) or a yeast cell (such as *Saccharomyces cerevisiae*), or into a baculovirus vector for expression in an insect cell. Such vectors comprise the regulatory elements necessary for  
30 expression of the DNA in the host cell, positioned in such a

manner as to permit expression of the DNA in the host cell. Such regulatory elements required for expression include promoter sequences, transcription initiation sequences and, optionally, enhancer sequences.

5           The *plPAC* polypeptide produced by gene expression in a recombinant procaryotic or eucyrotic system may be purified according to methods known in the art. In a preferred embodiment, a commercially available expression/secretion system can be used, whereby the  
10 recombinant protein is expressed and thereafter secreted from the host cell, to be easily purified from the surrounding medium. If expression/secretion vectors are not used, an alternative approach involves purifying the recombinant protein by affinity separation, such as by immunological  
15 interaction with antibodies that bind specifically to the recombinant protein. Such methods are commonly used by skilled practitioners.

          The *plPAC*-encoded polypeptides of the invention, prepared by the aforementioned methods, may be analyzed  
20 according to standard procedures.

          Polyclonal or monoclonal antibodies directed toward any of the peptides encoded by *plPAC* may be prepared according to standard methods. Monoclonal antibodies may be prepared according to general methods of Köhler and Milstein,  
25 following standard protocols.

### **C. Transgenic Plants**

          Transgenic plants expressing the *plPAC* gene can be generated using standard plant transformation methods known  
30 to those skilled in the art. These include, but are not

limited to, *Agrobacterium* vectors, PEG treatment of protoplasts, biolistic DNA delivery, UV laser microbeam, gemini virus vectors, calcium phosphate treatment of protoplasts, electroporation of isolated protoplasts, agitation of cell suspensions with microbeads coated with the transforming DNA, direct DNA uptake, liposome-mediated DNA uptake, and the like. Such methods have been published in the art. See, e.g., Methods for Plant Molecular Biology (Weissbach & Weissbach, eds., 1988); Methods in Plant Molecular Biology (Schuler & Zielinski, eds., 1989); Plant Molecular Biology Manual (Gelvin, Schilperoort, Verma, eds., 1993); and Methods in Plant Molecular Biology - A Laboratory Manual (Maliga, Klessig, Cashmore, Gruissem & Varner, eds., 1994).

The method of transformation depends upon the plant to be transformed. The biolistic DNA delivery method is useful for nuclear transformation. In another embodiment of the invention, *Agrobacterium* vectors are used to advantage for efficient transformation of plant nuclei.

In a preferred embodiment, the gene is introduced into plant nuclei in *Agrobacterium* binary vectors. Such vectors include, but are not limited to, BIN19 (Bevan, 1984, Nucleic Acid Res 12: 8711-8721) and derivatives thereof, the pBI vector series (Jefferson et al., 1987, PNAS 83:844751), and binary vectors pGA482 and pGA492 (An, 1986) and others (for review, see An, 1995, Methods Mol Biol 44:47-58). In preferred embodiments, the pPZP211 vector (Hajdukiewicz et al., 1994, PMB 25:989994) or PCGN7366 (Calgene, CA) are used. DNA constructs for transforming a selected plant comprise a coding sequence of interest operably linked to appropriate 5'

(e.g., promoters and translational regulatory sequences) and 3' regulatory sequences (e.g., terminators).

DNA constructs for transforming a selected plant comprise a coding sequence of interest operably linked to appropriate 5' (e.g., promoters and translational regulatory sequences) and 3' regulatory sequences (e.g., terminators). In a preferred embodiment, the coding region is placed under a powerful constitutive promoter, such as the Cauliflower Mosaic Virus (CaMV) 35S promoter or the figwort mosaic virus 35S promoter. Other constitutive promoters contemplated for use in the present invention include, but are not limited to: T-DNA mannopine synthetase, nopaline synthase (NOS) and octopine synthase (OCS) promoters.

Transgenic plants expressing a sense or antisense SDS coding sequence under an inducible promoter are also contemplated to be within the scope of the present invention. Inducible plant promoters include the tetracycline repressor/operator controlled promoter, the heat shock gene promoters, stress (e.g., wounding)-induced promoters, defense responsive gene promoters (e.g. phenylalanine ammonia lyase genes), wound induced gene promoters (e.g. hydroxyproline rich cell wall protein genes), chemically-inducible gene promoters (e.g., nitrate reductase genes, glucanase genes, chitinase genes, etc.) and dark-inducible gene promoters (e.g., asparagine synthetase gene) to name a few.

Tissue specific and development-specific promoters are also contemplated for use in the present invention. Examples of these included, but are not limited to: the ribulose biphosphate carboxylase (RuBisCo) small subunit gene promoters or chlorophyll a/b binding protein (CAB) gene

promoters for expression in photosynthetic tissue; the various seed storage protein gene promoters for expression in seeds; and the root-specific glutamine synthetase gene promoters where expression in roots is desired.

5           The coding region is also operably linked to an appropriate 3' regulatory sequence. In a preferred embodiment, the nopaline synthetase polyadenylation region (NOS) is used. Other useful 3' regulatory regions include, but are not limited to the octopine (OCS) polyadenylation  
10   region.

          Using an *Agrobacterium* binary vector system for transformation, the *plPAC* coding region, under control of a constitutive or inducible promoter as described above, is linked to a nuclear drug resistance marker, such as kanamycin  
15   resistance. *Agrobacterium*-mediated transformation of plant nuclei is accomplished according to the following procedure:

          (1) the gene is inserted into the selected *Agrobacterium* binary vector;

          (2) transformation is accomplished by co-  
20   cultivation of plant tissue (e.g., leaf discs) with a suspension of recombinant *Agrobacterium*, followed by incubation (e.g., two days) on growth medium in the absence of the drug used as the selective medium (see, e.g., Horsch et al. 1985, Cold Spring Harb Symp Quant Biol. 50:4337);

25           (3) plant tissue is then transferred onto the selective medium to identify transformed tissue; and

          (4) identified transformants are regenerated to intact plants.

          It should be recognized that the amount of  
30   expression, as well as the tissue specificity of expression

of the *plPAC* gene in transformed plants can vary depending on the position of their insertion into the nuclear genome. Such position effects are well known in the art. For this reason, several nuclear transformants should be regenerated and tested for expression of the transgene.

#### IV. Uses of *PlPAC* Nucleic Acids, Encoded Proteins and Antibodies

##### 1. *PlPAC* Nucleic Acids

*PlPAC* nucleic acids may be used for a variety of purposes in accordance with the present invention. The DNA, RNA, or fragments thereof may be used as probes to detect the presence of and/or expression of *plPAC* genes. Methods in which *plPAC* nucleic acids may be utilized as probes for such assays include, but are not limited to: (1) *in situ* hybridization; (2) Southern hybridization (3) northern hybridization; and (4) assorted amplification reactions such as polymerase chain reactions (PCR).

The *plPAC* nucleic acids of the invention may also be utilized as probes to identify related genes from other plant species. As is well known in the art and described above, hybridization stringencies may be adjusted to allow hybridization of nucleic acid probes with complementary sequences of varying degrees of homology. Thus, *plPAC* nucleic acids may be used to advantage to identify and characterize other genes of varying degrees of relation to the exemplary *ATPAC*, thereby enabling further characterization of this family of genes in plants.

Additionally, they may be used to identify genes encoding

proteins that interact with the P-glycoprotein encoded by *plPAC* (e.g., by the "interaction trap" technique).

Further, as described below, the genes or antisense molecules may be used to produce transgenic plants that have altered responses to herbicides and auxin.

## 2. PlPAC Proteins and Antibodies

Purified *plPAC*-encoded P-glycoproteins, or fragments thereof, may be used to produce polyclonal or monoclonal antibodies which also may serve as sensitive detection reagents for the presence and accumulation of plant P-glycoproteins in cultured plant cells or tissues and in intact plants.

Recombinant techniques enable expression of fusion proteins containing part or all of the *plPAC*-encoded protein. The full length protein or fragments of the protein may be used to advantage to generate an array of monoclonal or polyclonal antibodies specific for various epitopes of the protein, thereby providing even greater sensitivity for detection of the protein in cells or tissue.

Polyclonal or monoclonal antibodies immunologically specific for *plPAC*-encoded proteins may be used in a variety of assays designed to detect and quantitate the protein. Such assays include, but are not limited to: (1) flow cytometric analysis; (2) immunochemical localization in cultured cells or tissues; and (3) immunoblot analysis (e.g., dot blot, Western blot) of extracts from various cells and tissues.

Polyclonal or monoclonal antibodies that immunospecifically interact with one or more of the polypeptides encoded by *plPAC* can be utilized for identifying

and purifying such proteins. For example, antibodies may be utilized for affinity separation of proteins with which they immunospecifically interact. Antibodies may also be used to immunoprecipitate proteins from a sample containing a mixture  
5 of proteins and other biological molecules.

### 3. plPAC Transgenic Plants

Transgenic plants that over- or under- express plPAC can be used in a varied of agronomic and research  
10 applications. From the foregoing discussion, it can be seen that plPAC and its homologs, and transgenic plants containing them will be useful for improving stress resistance or tolerance in plants. This provides an avenue for developing marginal or toxic soil environments for crop production.  
15 Both over- and under-expressing plPAC transgenic plants have great utility in the research of herbicides and other xenobiotic compounds.

As discussed above and in greater detail in Example 1, the similarity between plant and mammalian *mdr* genes  
20 indicates that their functional aspects will also be conserved. Thus, plPAC is expected to play an important role in the exclusion of toxic metabolic or xenobiotic compounds from cells. The fact that plPAC also is inducible and appears to be preferentially expressed in roots, where  
25 contact with such compounds often occurs, makes plPAC particularly desirable for genetic engineering of plants to increase their tolerance to such compounds. Accordingly, plants engineered to overexpress the plPAC gene should be resistant to a wide range of chemicals, both intentionally  
30 applied as herbicides or unintentionally as wastes. Examples



of the kinds of xenobiotics that should be detoxified by the *plPAC* of the invention include, but are not limited to, hydrophobic (i.e., lipophilic) herbicides and other compounds, such as 3(3,4-dichlorophenyl)-1,1, dimethyl urea (also known as DCMU or Diuron, available from Sigma Chemical Co., St. Louis, MO) or other hydrophobic compounds that disrupt photosynthetic electron transport, as well as Metachlor (Ciba Geigy, Basel Switzerland), Taurocholate (Sigma Chemical Co.), Primisulfuron (Ciba Geigy), and IRL-1803.

As illustrated in Example 2, plant cells that over-express a *plPAC* gene have surprisingly higher growth rate with or without the xenobiotic compound Rhodamine 6G. It is contemplated that *plPAC* overexpression may be a generally useful way to increase plant and plant cell culture growth, even without the presence of xenobiotic compounds.

In addition to the above-mentioned features and advantages of transgenic plants that are altered in their expression of *plPAC*, these plants will also be altered in auxin transport. Through the use of developmental or tissue specific promoters, plants having a pre-determined alteration in auxin transport may be produced, providing agronomically or horticulturally beneficial features to such plants.

The following specific examples are provided to illustrate embodiments of the invention. They are not intended to limit the scope of the invention in any way.

#### EXAMPLE 1

##### Cloning and Analysis of a *PlPAC From Arabidopsis thaliana*

The *plPAC* of the present invention was identified by its up-regulation in response to a chloride ion channel blocker. *Brassica napus* plants were grown either in the presence or absence of 20  $\mu$ M 5-nitro-2-(3-phenylpropylamino) benzoic acid (NPPB). After five days, the roots of the seedlings were harvested and total RNA was extracted separately from the treated and untreated plants. From the total RNA preparations, poly (A)+ RNA was isolated and used as the starting material to create a cDNA subtraction library, using the CLONTECH PCR-SELECT™ cDNA Subtraction Kit and accompanying instructions (CLONTECH Laboratories, Inc., Palo Alto, CA).

Using the subtractive hybridization kit, a gene fragment was identified that was up-regulated in response to treatment of the plants with NPPB. This fragment (0.5 kb) was used to screen a cDNA library of *Arabidopsis thaliana*, from which a full-length cDNA clone was isolated. The nucleotide sequence of this cDNA clone, referred to as *ATPAC* (*Arabidopsis thaliana* putative anion channel) is set forth below as SEQ ID NO:1.

The 3.76 kb cDNA clone encodes a polypeptide 1,254 amino acids in length. The deduced amino acid sequence encoded by SEQ ID NO:1 is shown in Figure 1 as "atpac" (SEQ ID NO:2), in a lineup with the following sequences: (1) *hmdr1* (SEQ ID NO:3); (2) *mmdr1* (SEQ ID NO:4); (3) *hmdr3* (SEQ ID NO:5); (4) *mmdr2* (SEQ ID NO:6); (5) *atpgp1* (SEQ ID NO:7); and (6) *atpgp2* (SEQ ID NO:8). A consensus sequence (SEQ ID NO:9) is also shown.

A search of various sequence databases indicates that *ATPAC* is a new and distinct member of the *mdr* family of

ABC transporters. In none of the databases, including the EST collection, does an exact match exist. The ABC transporter family is very large, consisting of at least two sub-groups, *mrp* and homologs and *mdr* and homologs. The only examples of plant *mdr*-like genes are *atpgp1* and *atpgp2* from *A. thaliana* and two homologs from potato and barley, respectively. Though the *atpgp1* and *atpgp2* genes are similar to *ATPAC*, they are only 51 and 50% identical, respectively, indicating that *ATPAC* is a distinct gene by comparison.

Sequence homology with the potato and barley *mdr*-like genes is even more divergent. Another difference between the *agpgp1* gene and the *ATPAC* gene is their respective preferential expression in inflorescens and roots, respectively.

## EXAMPLE 2

### Effect of *ATPAC* Expression in Bacterial Cells on Their Ability to Detoxify Rhodamine 6G

The compound Rhodamine 6G is a well known substrate of mammalian p-glycoproteins (Kolaczowski et al., J. Biol. Chem. 271: 31543-31548, 1996). The ability of a cell to detoxify the compound is indicative of activity of p-glycoproteins. A bacterial cell line was transformed with an expression vector comprising *ATPAC*. The growth rate of transformed and non-transformed cells was then measured, in the presence or absence of Rhodamine 6G. Results are shown in Figure 2. As can be seen, *ATPAC*-expressing cells grown in the absence of the drug had the best growth rate. Moreover, even in the presence of the drug, the cells grew more quickly than non-transformed cells in the presence or absence of

Rhodamine 6G. These results demonstrate that *ATPAC* encodes a functional and robust p-glycoprotein.

**Example 3**  
**Transgenic Plants that Overexpress**  
**and Underexpress ATPAC**

**Transformation construct.** The *Agrobacterium* binary vector pPZP211 (Hajdukiewicz et al., 1994 Plant Mol. Biol. 25:989994) was digested with *EcoRI* and *SmaI*, and selfligated. This molecule was named pPZP211'. The *Agrobacterium* binary vector pCGN7366 (Calgene, CA) was digested with *XhoI* and cloned in *SalI* digested pPZP211'. We named this binary vector pPZPPCGN. The 3.8 kb full-length *ATPAC* cDNA was cloned into the pGH19 vector. After digestion with *SmaI* (in the multiple cloning site upstream) and *EcoRI*, a 3.1 kb cDNA fragment was cut out. This *SmaI**EcoRI* 3.1 kb fragment was cloned into the *SmaI*/*EcoRI* site of pPZPPCGN. The rest of *ATPAC* gene was amplified using polymerase chain reaction to have translationally fused HAtag at its 3'terminal. After ligating *EcoRI* linkers to the ends of the resulting PCR product, the 0.7 kb fragment was cloned into the *EcoRI* site of the *SmaI**EcoRI* 3.1 kb *ATPAC* fragment in pPZP-pCGN. The final construct was named pATPACOE.

**Plant transformation.** pATPACOE was introduced into *Agrobacterium tumefaciens* strain by a direct transformation method. *Agrobacterium*-mediated transformation was performed using vacuum infiltration (Bechtold et al., 1993, . CR Acad. Sci. [III] 316: 11941199.)

T1 plants which survived on kanamycincontaining plates were selected, transplanted into soil and grown to set

T2 seed. T3 seeds were collected from kanamycinresistant T2 plants. T3 plants which showed 100% kanamycinresistance were selected and were considered homozygous for the transgene.

**Antisense Plants.** The full length cDNA in  
5 pBluescript SK() vector (Stratagene, CA) is digested with  
*EcoRI* (there is a cleavage site in the upstream polylinker)  
and *SspI*. The resulting 1.3 Kb fragment representing a 5'  
portion of the *AtPAC* cDNA was cloned into the aforementioned  
pPZPPCGN, which had been digested with *EcoRI/SmaI*, ensuring  
10 that this fragment of the cDNA was inserted in the antisense  
orientation. This construct was named *pATPACAE*. *pATPACAE*  
was introduced into *Arabidopsis* plants by *Agrobacterium*  
transformation, as described above.

**Knock-out Plants.** The method of Krysan et al  
15 (1996, PNAS 93:8145, incorporated by reference herein) was  
followed using the following primers:

Genespecific primers:

*AtpacF*: CACTGCTCAATGATCTCGTTTTCTCACTA (SEQ ID NO:11)

*AtpacR*: CTTGAATCACACCAATGCAATCAACACCTC (SEQ ID NO:12)

20 Primers for TDNA left boarder:

JL202: CATT TTTATAATAACGCTGCGGACATCTAC (SEQ ID NO:13)

JL270: TTTCTCCATATTGACCATCATACTCATTG (SEQ ID NO:14)

T-DNA insertion mutants were isolated by PCR-based  
25 screening of DNA pools. Two alleles of *ATPAC* and one *ATPGPI*  
allele were isolated. At the seedling stage, both alleles of  
*ATPAC* displayed down-folded, or epinastic cotyledons and  
abnormally angled cotyledon petioles. The petioles of the  
cotyledons and first true leaves in *ATPAC* were shorter than  
30 in wild type plants. As adult plants, the rosette leaves

were somewhat curled and wrinkled along the margin. Bolting of the inflorescence stem was delayed by 2.8 days on average, relative to wild-type. The bolt grew more slowly than wild type after starting, though the wild type length was ultimately reached. These phenotypes coincided with the sites of expression indicated by GUS staining (as described herein). None of these phenotypes were present in plants transformed with a genomic fragment containing the wild type *ATPAC* promoter and coding sequence ("*atpac1-1*"). This mutant did not display any overt phenotype as a seedling or an adult plant. Double mutants were constructed by crossing *atpac1-1* and *atpgp1-1* plants. F1 individuals appeared wild type and were permitted to self pollinate. Approximately one in sixteen of the F2 seedlings displayed extremely down-curved cotyledons when grown in the light and also displayed shorter, wavy hypocotyls when grown in the dark. PCR analysis confirmed that these seedlings were homozygous double mutants. The adult double mutant plants were severely stunted in growth. Primary inflorescence stems of the double mutants and floral pedicels were also wavy in appearance, indicating that the direction of growth periodically changed during the elongation phase of these organs. After 72 hours of growth, the double mutant produced abundant secondary inflorescence stems, indicating a large reduction in apical dominance. Fertility of the flowers in the double mutants was poor, due to improper elongation of the stamen filament. Hand-pollinating flowers of the doubly homozygous mutant plants produced double-mutant seeds that were both viable and normal in appearance.

#### Example 4

##### Effect of Auxin (IAA) on ATPAC Expression

###### **Expression in Yeast and Xenopus Oocytes. ATPAC**

5 cDNA was expressed in wild type yeast as well as in yeasts  
lacking seven ABC transporters (as described by Decottignies  
et al., (1998) J. Biol. Chem. 273:12612-22) in order to create  
a heterologous system for studying function of the  
transporter. The yeasts were exposed to toxic compounds that  
10 are known substrates for human MDR1. ATPAC did not confer  
any measurable resistance to the toxic substrates. Further,  
there was no evidence of a drug-pumping role for ATPAC.

In order to examine whether ATPAC functions as an  
anion channel or a regulator of an anion channel,  
15 complementary RNA made from an ATPAC cDNA template was  
injected into Xenopus oocytes to produce a heterologous  
expression system amenable to electrophysiology. No currents  
associated with ATPAC were observed by two electrode voltage  
clamping.

20 Treatment of wild type seedlings with 2,4-D or high  
concentrations of auxin result in plants having strongly  
curved organs.

###### **ATPAC Binds the Auxin Transport Inhibitor**

**naphthylphthalamic acid (NPA).** Yeast expressing ATPAC was  
25 assayed for NPA binding. NPA bound tightly and specifically  
to ATPAC-expressing yeast, but not to control yeast. Bound  
NPA was not displaced by benzoic acid, a related hydrophobic  
organic acid. Proteins that bind NPA and influence auxin  
transport are considered important regulators of hormone  
30 action and therefore, of plant development. The fact that

NPA binds to ATPAC and that *atpac* knockout mutants can be phenocopied by auxin application suggests that ATPAC is an important component of the auxin transport and distribution machinery.

5                   **Effect of ATPAC and AtPGP1 on Auxin Transport.**

Three different auxin transport assays of the Ws wild type, ATPAC-1, ATPAC-2 and *atpgp1* were performed. The first measured the basipetal movement of auxin in seedlings as described by Murphy et al. (2000) *Planta* 211:315-324. A 0.1  
10    μl microdroplet of radioactive auxin was placed on the apex of a light-grown seedling. Four hours later, the amount of radioactivity collected on moist filter paper that contacted the root tip was determined. Polar auxin transport measured in this manner was severely reduced in both alleles of ATPAC  
15    and especially in the double mutant, but not in the *atpgp1* mutant.

The second assay measured the basipetal transport of radioactive auxin through etiolated seedlings inverted in a reservoir containing radioactive auxin (Garbers et al.  
20    (1996) *EMBO J.* 15:2115-2124). The results were similar to those with the first assay described above.

The third assay measured basipetal transport of auxin in the inflorescence stem in a method described by Ruegger et al (1997) *Plant Cell* 9:745-57. A segment of  
25    inflorescence stem was excised and immersed apical-end down in a tube containing a small volume of radioactive auxin. At a later time, a piece of tissue was excised from the basal end of the segment and counted in a scintillation counter. Seedlings having ATPAC mutations clearly displayed defective  
30    auxin transport profiles. Further, the *atpgp1* mutation



significantly impaired transport in tissue segments taken from the lower portion of the inflorescence. This is indicative of gradients of function of at least two MDR-like gene products along the inflorescence axis.

5

While certain of the preferred embodiments of the present invention have been described and specifically exemplified above, it is not intended that the invention be limited to such embodiments. Various modifications may be made thereto without departing from the scope and spirit of the present invention, as set forth in the following claims.

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209